

Use this replacement page in all versions of the *Goat Project and Record Book* published before November 2025. This feed label and answers are up-to-date.

Activity 2

Read the following and answer the questions.

Goat Starter

MEDICATED

Goats in confinement except reproducing or lactating goats
Complete Feed
Prevention of coccidiosis caused by *Eimeria crandallis*,
E. christenseni, and *E. ninakohlyakimovae*.

Active Drug Ingredient

Monensin 20 gm/ton

Guaranteed Analysis

Crude Protein, min % 18.0	Salt (NaCl), min % 0.1
Equiv. Cr Prot from NPN, max % 1.0	Salt (NaCl), max % 0.6
Crude Fat, min % 3.7	Sodium (Na), min % 0.01
Crude Fiber, max % 11.6	Sodium (Na), max % 0.5
ADF, max % 14.3	Copper (Cu), min ppm .. 10.00
Calcium (Ca), min % 0.9	Copper (Cu), max ppm .. 30.00
Calcium (Ca), max % 1.4	Selenium (Se), min ppm .. 0.30
Phosphorus (P), min % 0.4	Vitamin A, min IU/lb 1,500

Ingredient Statement

Forage Products, Grain Products, Roughage Products, Plant Protein Products, Processed Grain By-Products, Molasses Products, Choice White Grease, Ammonium Chloride, Dicalcium Phosphate, Limestone Ground, Bentonite, Zinc Amino Acid Complex, Manganese Amino Acid Complex, Copper Amino Acid Complex, Cobalt Glucoheptonate, Vitamin A Supplement, Sodium Selenite, Salt, Vitamin E Supplement

Directions For Use

Feed continuously and only to goats in confinement.
Feed to goat kids 1-8 weeks of age. Provide clean, fresh water at all times.

WARNING. Do not feed to lactating goats.

CAUTION. USE AS DIRECTED. Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated goat feeds are safe for use in goats only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result.

Manufactured By:
Skillathon Feeds

Net Wt. 50 lbs

1. At what growth stage of development should this ration be fed?

2. What is the main ingredient in this feed?

3. What is the minimum crude protein level?

4. What is the minimum crude fat level of this diet?

5. Is this a medicated feed?

6. What is the active drug ingredient?

7. Is there a withdrawal time for this feed?

ANSWERS:
1. 1-8 weeks of age
2. Forage products
3. 18%
4. 3.7%
5. yes
6. monensin
7. no

